

High-Level Research Summaries For Key Issues

ISSUES COVERED

Trends in Forest
Gain/Loss

Biodiversity

Forest and Water

Forest Carbon

Forest Gain/Loss – Southeastern U.S.

- Harvesting rates in the southeastern U.S. have slightly declined and there continues to be a surplus of growth compared to harvest (NCASI 2022¹).
- The southeastern U.S. continues to grow far more timber than is harvested. Annual growth amounts to 681.6 million green tons, while harvest totals 390.8 million tons (USDA 2022²).
- It is important to understand that the primary driver of forest loss in the southeastern U.S. is conversion due to urbanization (Olson 2020³).



Biodiversity – Southeastern U.S.

- Research has demonstrated that active forest management within managed pine landscapes contributes to conservation of biological diversity (e.g., Loehle et al. 2006⁴; Miller et al. 2009⁵; Verschuyt et al. 2011⁶; Iglay et al. 2012⁷, 2014⁸, 2018⁹; King and Schlossbert 2014¹⁰; Bender et al. 2015¹¹; Demarais et al. 2017¹²; Parrish et al. 2017¹³; Levy et al. 2025¹⁴).
- Different species require different forest conditions; therefore, there is no one set of forest management recommendations that will benefit all species in a single stand (e.g., Guldin et al. 2007¹⁵; Favorito et al. 2023¹⁶).
- At the landscape scale, forest management can provide for the

changing needs of species throughout the year (e.g., Edwards et al. 2004¹⁷; Miller and Conner 2005¹⁸; Brooks 2009¹⁹; Bender et al. 2015²⁰; Homyack et al. 2016²¹; Guzy et al. 2019a²², 2019b²³; Perea et al.

Water from Forests and Best Management Practices (BMPs)

- State and private forests contribute 370 billion m³ yr⁻¹ to the surface water supply with approximately 55 million people in the southeastern US deriving some portion of their drinking water from private forests (Liu et al. 2020²⁵).
- Forestry BMPs, when properly applied, are highly effective at reducing erosion and the potential for sediment delivery to waterbodies in the southeastern US (Cristan et al. 2016²⁶; Fielding et al. 2022²⁷; Hawks et al. 2022²⁸; Hawks et al. 2023²⁹).
- Application of forestry BMPs coupled with state monitoring programs and participation in forest certification programs, that require routine third-party audits, provide assurance to federal and state agencies that BMPs protect aquatic resources and species (Warrington et al. 2017³⁰; Schilling et al. 2021³¹).

Forest Carbon

- Forests remove CO₂ from the atmosphere and store it in live trees, dead wood, and harvested wood products; therefore, sustainably managed forests play a key role in mitigating effects of climate change (Nabuurs et al. 2007³²).
- While it is true that mature and old growth forests store more C than younger forests, younger forests sequester C at a much faster rate (Gray et al. 2016³³).
- Forest growth trajectories show more rapid growth at young ages than at older ones, therefore maximizing C storage can be best achieved at harvest rotations near the culmination of mean annual increment (peak of average annual growth; Diaz et al. 2018³⁴).
- While reduced harvest levels may lead to an increase in forest C stocks, it may also lead to increased use of substitute products that are accompanied by much higher emissions from production and use (Churkina et al. 2020³⁵; NCASI 2020³⁶).



CONCEPT CONNECTIONS

- Younger forests sequester carbon at a faster rate than older forests and are important for diversity of wildlife species, including some in decline.
- Active forest management, including implementing BMPs, at a landscape scale, maintains a diversity of forest and cover types and conditions to support biodiversity, water quality, and carbon sequestration and storage.
- Economic return on forests encourages maintaining ecosystem services and sustainable wood fiber.



REFERENCES

1. NCASI. 2022. Trends in forest harvest, regeneration, and management in the southeastern United States as related to biomass feedstock. Briefing Note. Cary, NC: National Council for Air and Stream Improvement, Inc.
2. USDA. 2022. Forest Inventory and Analysis Program, Forest Inventory EVALIDator, Version 1.8.0.01. Department of Agriculture, Forest Service, Northern Research Station, St. Paul, MN. <https://apps.fs.usda.gov/fiadb-api/evaluator>
3. Olson, R.L. 2020. Demographics as a driver of change in the U.S. forest sector. In Dockry, M.J., Bengston, D.N., and Westphal, L.M. (comps.) *Drivers of change in U.S. forests and forestry over the next 20 years*. General Technical Report NRS-P-197. Madison, WI: U.S. Department of Agriculture, Forest Service, Northern Research Station: 59–67. <https://doi.org/10.2737/NRS-GTR-P-197-paper6>
4. Loehle, C., Van Deusen, P., Wigley, T.B., Mitchell, M.S., Rutzmoser, S.H., Aggett, J., Beebe, J.A., and Smith, M.L. 2006. A method for landscape analysis of forestry guidelines using bird habitat models and the Habplan harvest scheduler. *Forest Ecology and Management* 232:56-67. <https://doi.org/10.1016/j.foreco.2006.05.040>.
5. Miller, D.A., Wigley, T.B., and Miller, K.V. 2009. Managed forests and conservation of terrestrial biodiversity in the southern United States. *Journal of Forestry* 107:197-203. <https://doi.org/10.1093/jof/107.4.197>.
6. Verschuy, J.P., Riffell, S., Miller, D.A., and Wigley, T.B. 2011. Biodiversity response to intensive biomass production from forest thinning in North American forests - a meta-analysis. *Forest Ecology and Management* 261:221-232. <https://doi.org/10.1016/j.foreco.2010.10.010>.
7. Iglay, R.B., Demarais, S., Wigley, T.B., and Miller, D.A. 2012. Bird community dynamics and vegetation relationships among stand establishment practices in intensively managed pine stands. *Forest Ecology and Management* 283:1-9. <https://doi.org/10.1016/j.foreco.2012.07.008>.
8. Iglay, R.B., Leopold, B.D., and Miller, D.A. 2014. Vegetation responses to fire and herbicide in intensively managed, mid-rotation pine. *Forest Ecology and Management* 328:69-78. <https://doi.org/10.1016/j.foreco.2014.05.029>.
9. Iglay, R.B., Greene, R.E., Leopold, B.D., and Miller, D.A. 2018. Bird conservation potential of fire and herbicide treatments in thinned pine stands. *Forest Ecology and Management* 409:267-275. <https://doi.org/10.1016/j.foreco.2017.11.021>.
10. King, D.I., and Schlossberg, S. 2014. Synthesis of the conservation value of the early-successional stage in forests of eastern North America. *Forest Ecology and Management* 324:186-195. <https://doi.org/10.1016/j.foreco.2013.12.001>.
11. Bender, M.J., Castleberry, S.B., Miller, D.A., and Wigley, T.B. 2015. Site occupancy of foraging bats on landscapes of managed pine forest. *Forest Ecology and Management* 336:1-10. <https://doi.org/10.1016/j.foreco.2014.10.004>.
12. Demarais, S., Verschuy, J.P., Roloff, G.J., Miller, D.A., and Wigley, T.B. 2017. Tamm Review: Terrestrial vertebrate biodiversity and intensive forest management in the U.S. *Forest Ecology and Management* 385:308-330. <https://doi.org/10.1016/j.foreco.2016.10.006>.
13. Parrish, M.C., Demarais, S., Wigley, T.B., Jones, P.D., Ezell, A.W., and Riffell, S.K. 2017b. Breeding bird communities associated with land cover in intensively managed pine forests of the southeastern U.S. *Forest Ecology and Management* 406:112-124. <https://doi.org/10.1016/j.foreco.2017.09.063>.

14. Levy, Fen S., J. Michael Reed, Peter S. McKinley, John S. Gunn, Kelsi Anderson, and John M. Hagan. 2025. Increased bird abundances over 30 years in an extensive commercial forest landscape." *Biological Conservation* 302: 110934.
15. Guldin, J.M., Emmingham, W., Carter, S.A., and Saugey, D. 2007. Silvicultural practices and management of habitat for bats. 177-205 in Lacki, M.J., Hayes, J.P., and Kurta, A. (eds.). *Bats in Forests: Conservation and Management*. Baltimore, MD: John Hopkins University Press.
16. Favorito, C. M., B. F. Barnes, E. L. Briggs, C. C. Fortuin, D. U. Greene, A. L. Larsen-Gray, J. A. Martin, E. McCarty, and K. J. K. Gandhi. 2023. Stand stage affects wild bee communities in working pine forests. *Forest Ecology and Management* 545: 121247.
17. Edwards, S.L., Demarais, S., Watkins, B., and Strickland, B.K. 2004. White-tailed deer forage production in managed and unmanaged pine stands and summer food plots in Mississippi. *Wildlife Society Bulletin* 32:739-745. [https://doi.org/10.2193/0091-7648\(2004\)032\[0739:WDFPIM\]2.0.CO;2](https://doi.org/10.2193/0091-7648(2004)032[0739:WDFPIM]2.0.CO;2).
18. Miller, D.A., and Conner, L.M. 2005. Seasonal and annual home ranges of female eastern wild turkeys in a managed pine landscape in Mississippi. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 59:89-99.
19. Brooks, R.T. 2009. Habitat-associated and temporal patterns of bat activity in a diverse forest landscape of southern New England, USA. *Biodiversity Conservation* 18:529-545. <https://doi.org/10.1007/s10531-008-9518-x>.
20. Bender, M.J., Castleberry, S.B., Miller, D.A., and Wigley, T.B. 2015. Site occupancy of foraging bats on landscapes of managed pine forest. *Forest Ecology and Management* 336:1-10. <https://doi.org/10.1016/j.foreco.2014.10.004>.
21. Homyack, J.A., O'Bryan, C.J., Thornton, J.E., and Baldwin, R.F. 2016. Community occupancy of herpetofauna in roadside ditches in a managed pine landscape. *Forest Ecology and Management* 361:346-357. <https://doi.org/10.1016/j.foreco.2015.11.037>.
22. Perea, Santiago, Amanda Vicente-Santos, Angela L. Larsen-Gray, Kamal JK Gandhi, Daniel U. Greene, Brittany F. Barnes, and Steven B. Castleberry. 2025. Disentangling winter relationships: Bat responses to forest stand structure, environmental conditions, and prey composition. *Forest Ecology and Management* 578: 122484.
23. Guzy, J., Halloran, K., Homyack, J., Thornton-Frost, J., and Willson, J. 2019a. Differential responses of amphibian and reptile assemblages to size of riparian buffers within managed forests. *Ecological Applications* 29:e01995. <https://doi.org/10.1002/eap.1995>.
24. Guzy, J., Halloran, K., Homyack, J., and Willson, J.D. 2019b. Influence of riparian buffers and habitat characteristics on salamander assemblages in headwater streams within managed forests. *Forest Ecology and Management* 432:868-883. <https://doi.org/10.1016/j.foreco.2018.10.006>.
25. Liu, N., P.V. Caldwell, G.R. Dobbs, C.F. Miniati, P.V. Bolstad, S.A.C. Nelson, and G. Sun. 2021. Forest lands dominate drinking water supply in the conterminous United States. *Environmental Research Letters* 16. <https://doi.org/10.1088/1748-9326/ac09b0>.
26. Cristan, R., Aust, W.M., Bolding, M.C., Barrett, S.M., Munsell, J.F., Schilling, E., 2016. Effectiveness of forestry best management practices in the United States: Literature review. *Forest Ecology Management*. 360, 133–151. <https://doi.org/10.1016/j.Foreco.2015.10.025>.
27. Fielding, J.A.H., Hawks, B.S., Aust, W.M., Bolding, M.C., Barrett, S.M., 2022. Estimated erosion from clearcut timber harvests in the southeastern United States. *Forest Science*. 68, 334–342. <https://doi.org/10.1093/forsci/fx013>.

28. Hawks, B.S., Aust, W.M., Bolding, M.C., Barrett, S.M., Schilling, E., Fielding, J.A.H., 2022. Linkages between forestry best management practices and erosion in the southeastern U.S. *Journal of Environmental Management*. 305, 114411. <https://doi.org/10.1016/j.jenvman.2021.114411>.
29. Hawks, B., M.C. Bolding, W.M. Aust, S.M. Barrett, E.B. Schilling, S.P. Prisley. 2023. Increased levels of forestry best management practices reduce sediment delivery from Piedmont and Upper Coastal Plain clearcut harvests and access features, southeastern states, USA. *Forest Ecology and Management*. <https://doi.org/10.1016/j.foreco.2022.120697>.
30. Warrington, B.M., W.M. Aust, S.M. Barrett, W.M. Ford, C.A. Dolloff, E.B. Schilling, T.B. Wigley, and M. C. Bolding. 2017. Forestry best management practices relationships with aquatic and riparian fauna: A review. *Forests* 8(9):331. doi:10.3390/f8090331.
31. Schilling, E.B., A.L. Larsen-Gray, and D.A. Miller. 2021. Forestry best management practices and conservation of aquatic systems in the southeastern United States. *Water*: 13(19): 2611.
32. Nabuurs, G.J., Masera, O., Andrasko, K., Benitez-Ponce, P., Boer, R., Dutschke, M., Elsiddig, E., Ford-Robertson, J., Frumhoff, P., Karjalainen, T., Krankina, O., Kurz, W.A., Matsumoto, M., Oyhantcabal, W., Ravindranath, N.H., Sanz Sanchez, M.J., and Zhang, X. 2007. Forestry. In Metz, O.R., Davidson, P.R. Bosch, R. Dave, L.A. Meyer (eds). *Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change [B]*. Cambridge, United Kingdom and New York, NY: Cambridge University Press.
33. Gray, A.N., Whittier, T.R., and Harmon, M.E. 2016. Carbon stocks and accumulation rates in Pacific Northwest forests: role of stand age, plant community, and productivity. *Ecosphere* 7:e01224. <https://doi.org/10.1002/ecs2.1224>.
34. Diaz, D.D., Loreno, S., Ettl, G.J., and Davies, B. 2018. Tradeoffs in timber, carbon, and cash flow under alternative management systems for Douglas-Fir in the Pacific Northwest. *Forests* 9:447. <https://doi.org/10.3390/f9080447>.
35. Churkina, G., Organschi, A., Reyer, C.P.O., Ruff, A., Vinke, K., Liu, Z., Reck, B.K., Graedel, T.E., and Schellnhuber, H.J. 2020. Buildings as a global carbon sink. *Nature Sustainability* 3:269-276. <https://doi.org/10.1038/s41893-019-0462-4>.
36. NCASI. 2020. NCASI review of carbon implications of proforestation. NCASI Review and Response. Cary, NC: National Council for Air and Stream Improvement, Inc.